

GENETIC TESTING: LUNG DISORDERS

OVERVIEW

One of the most common forms of inherited lung disorders is alpha-1 antitrypsin deficiency (AATD). AATD is an autosomal recessive genetic disorder that results in decreased production of the alpha-1 antitrypsin (AAT) protein, or production of abnormal types of the protein that are functionally deficient. Individuals with AATD have an increased risk to develop lung and liver disease. Genetic testing to diagnose AATD aids in directing proper treatment and identifying at-risk family members.

POLICY REFERENCE TABLE

The tests and associated laboratories and CPT codes contained within this document serve only as examples to help users navigate claims and corresponding coverage criteria; as such, they are not comprehensive and are not a guarantee of coverage or non-coverage. Please see the [Concert Genetics Platform](#) for a comprehensive list of registered tests.

Coverage Criteria Sections	Example Tests (Labs)	Common CPT Codes	Common ICD Codes	Ref
Alpha-1 Antitrypsin Deficiency				
SERPINA1 Common Variant Analysis or Sequencing and/or Deletion/Duplication Analysis	Alpha-1 Antitrypsin (AAT) Mutation Analysis (Quest Diagnostics)	81332	E88.01	1
	<i>SERPINA1</i> Full Gene Sequencing and Deletion/Duplication (Invitae)	81479		
Other Covered Lung Disorders				

Other Covered Lung Disorders	See list below	81400-81408		2, 3, 4
--	----------------	-------------	--	---------

OTHER RELATED POLICIES

This policy document provides coverage criteria for Genetic Testing for Lung Disorders. Please refer to:

- **Genetic Testing: Multisystem Inherited Disorders, Intellectual Disability, and Developmental Delay** for coverage criteria related to diagnostic testing for cystic fibrosis and other multisystem inherited disorders.
- **Genetic Testing: General Approach to Genetic and Molecular Testing** for coverage criteria related to genetic testing for lung disorders and disease that are not specifically discussed in this or another non-general policy.

[back to top](#)

COVERAGE CRITERIA

ALPHA-1 ANTITRYPSIN DEFICIENCY

SERPINA1 Common Variant Analysis or Sequencing and/or Deletion/Duplication Analysis

- I. **SERPINA1** common variant analysis (81332) or sequencing and/or deletion/duplication analysis (81479) to establish a diagnosis of alpha-1 antitrypsin (AAT) deficiency is considered **medically necessary** when:
 - A. The member has any of the following:
 1. Abnormally low (less than 120 mg/dL) or borderline (90-140 mg/dL) alpha-1 antitrypsin levels (as measured by nephelometry), **OR**
 2. Early-onset emphysema (45 years of age or younger), **OR**
 3. Emphysema in the absence of additional risk factor (e.g., smoking, occupational dust exposure), **OR**

4. Emphysema with prominent basilar hyperlucency, **OR**
 5. Otherwise unexplained liver disease, **OR**
 6. Necrotizing panniculitis, **OR**
 7. C-ANCA positive vasculitis (i.e., granulomatosis with polyangiitis),
OR
 8. Bronchiectasis without evident etiology, **OR**
 9. A sibling with known AAT deficiency.
- II. *SERPINA1* common variant analysis (81332) or sequencing and/or deletion/duplication analysis (81479) to establish a diagnosis of alpha-1 antitrypsin deficiency is considered **investigational** for all other indications.

[back to top](#)

OTHER COVERED LUNG DISORDERS

The following is a list of conditions that have a known genetic association. Due to their relative rareness, it may be appropriate to cover these genetic tests to establish or confirm a diagnosis.

- I. Genetic testing to establish or confirm one of the following genetic lung disorders to guide management is considered **medically necessary** when the member demonstrates clinical features* consistent with the disorder (the list is not meant to be comprehensive, see II below):
 - A. [Familial Pulmonary Fibrosis](#)
 - B. [Primary Ciliary Dyskinesia](#)
 - C. Pulmonary lymphangiomyomatosis (LAM)
 - D. Pulmonary alveolar proteinosis (PAP)
- II. Genetic testing to establish or confirm the diagnosis of all other lung disorders not specifically discussed within this or another medical policy will be evaluated by the criteria outlined in *General Approach to Genetic and Molecular Testing* (see policy for coverage criteria).

*Clinical features for a specific disorder may be outlined in resources such as [GeneReviews](#), [OMIM](#), [National Library of Medicine](#), [Genetics Home Reference](#), or other scholarly source.

[back to top](#)

BACKGROUND AND RATIONALE

***SERPINA1* Common Variant Analysis or Sequencing and/or Deletion/Duplication Analysis**

American Thoracic Society and European Respiratory Society

The American Thoracic Society and European Respiratory Society published a joint statement on the diagnosis and management of individuals with alpha-1 antitrypsin deficiency (2003) which provided recommendations for diagnostic testing.

A normal range of plasma alpha-1 antitrypsin (measured via nephelometry) is 83/120 - 200/220 mg/dL. Individuals with borderline normal levels of plasma alpha-1 antitrypsin (90-140 mg/dL) or with abnormally low levels (below 120 mg/dL) should be evaluated for alpha-1 antitrypsin deficiency. (p. 826 and 827)

“The following features should prompt suspicion by physicians that their patient may be more likely to have AAT deficiency:

- Early-onset emphysema (age of 45 years or less)
- Emphysema in the absence of a recognized risk factor (smoking, occupational dust exposure, etc.)
- Emphysema with prominent basilar hyperlucency
- Otherwise unexplained liver disease
- Necrotizing panniculitis
- Anti-proteinase 3-positive vasculitis (C-ANCA [anti-neutrophil cytoplasmic antibody]-positive vasculitis)
- Family history of any of the following: emphysema, bronchiectasis, liver disease, or panniculitis
- Bronchiectasis without evident etiology...” (p. 820)

The statement also recommended that individuals with a sibling with AAT deficiency should also be offered genetic testing. (p. 827)

[back to top](#)

REFERENCES

1. American Thoracic Society; European Respiratory Society. American Thoracic Society/European Respiratory Society statement: standards for the diagnosis and management of individuals with alpha-1 antitrypsin deficiency. *Am J Respir Crit Care Med.* 2003;168(7):818-900. doi:10.1164/rccm.168.7.818
2. Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1116/>
3. Online Mendelian Inheritance in Man, OMIM®. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD). World Wide Web URL: <https://omim.org/>
4. MedlinePlus [Internet]. Bethesda (MD): National Library of Medicine (US). Available from: <https://medlineplus.gov/genetics/>.

[back to top](#)