

*Evolent	
Clinical guidelines CARDIAC RESYNCHRONIZATION THERAPY (CRT)	Original Date: February 2013
CPT Codes: 33221, 33224, 33225, 33231	Last Revised Date: April 2023
Guideline Number: Evolent_CG_320	Implementation Date: January 2024

GENERAL INFORMATION

- *It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.*
- *Where a specific clinical indication is not directly addressed in this guideline, medical necessity determination will be made based on widely accepted standard of care criteria. These criteria are supported by evidence-based or peer-reviewed sources such as medical literature, societal guidelines and state/national recommendations.*

INDICATIONS FOR CARDIAC RESYNCHRONIZATION THERAPY (CRT)¹⁻⁸

Indications for CRT for patients are based upon LV ejection fraction (LVEF), QRS duration, New York Heart Association (NYHA) functional class (presence or absence of symptoms) and need for ventricular pacing regardless of etiology (ischemic or non-ischemic cardiomyopathy). The beneficial effects of CRT have been extensively proven in patients with NYHA class II, III, and IV; there is limited evidence of CRT benefit in patients with NYHA functional class I. Other special situations, such as patients with atrial fibrillation or who require an upgrade from a conventional pacing or ICD system, will be addressed below as well.

Patients with cardiomyopathy on GDMT for 3 months or on GDMT and 40 days after MI; or with implantation of pacing or defibrillation device for special indications

CRT-D Indications by NYHA Heart Failure Class (see full definitions further below in document). See [Background](#) for Algorithm for CRT Indications/Recommendations in patients with cardiomyopathy or HFrEF chart.

- Class II- Ambulatory IV
 - LVEF ≤ 35%, QRS ≥ 120ms, LBBB, Sinus Rhythm
 - LVEF ≤ 35%, QRS ≥ 150ms, non-LBBB, Sinus Rhythm

Special Situations

- Independent/Regardless of NYHA Heart Failure Class
 - Patients who have an indication for ventricular pacing and high degree AV block or are expected to be paced more than 40% of the time; this includes patients with Atrial fibrillation
- Atrial fibrillation and LVEF \leq 35% if:
 - Patient requires ventricular pacing or otherwise meets CRT criteria; **AND**
 - AV nodal ablation or pharmacologic rate control will allow nearly 100% ventricular pacing with CRT
 - For patients with atrial fibrillation and LVEF \leq 50%, if a rhythm control strategy fails and ventricular rates remain rapid despite medical therapy, atrioventricular nodal ablation with implantation of a CRT device is reasonable
- In patients with nonobstructive HCM who have NYHA class II to IV heart failure with LBBB, LVEF $<$ 50%, CRT therapy for symptom reduction is reasonable

NOT Indicated for Cardiac Resynchronization Therapy (CRT)

- NYHA class I and non-LBBB pattern with QRS duration $<$ 150 ms,³ except as in Special Situations section above
- Inotrope-dependent patients who have a higher risk need for cardiac transplant and LVAD support, are less likely to benefit from CRT
- Comorbidities and/or frailty expected to limit survival with good functional capacity to $<$ 1 year
- Active bloodstream infection
- Reversible causes are present such as toxic-, metabolic- or tachycardic-mediated cardiomyopathy, would require reassessment once the situation is corrected
- CRT has not been studied in ATTR-CM with HFrEF

Indications for CRT in Adult Congenital Heart Disease⁹⁻¹¹

Systemic LV

- Systemic LV EF \leq 35%, sinus rhythm, wide QRS complex \geq 130 ms NYHA function Class II—IV

Any Systemic V

- Systemic ventricle any EF (not restricted to \leq 35%), intrinsic narrow QRS complex, NYHA function Class I—IV and are undergoing new device placement or replacement with anticipated requirement for significant ($>$ 40%) ventricular pacing.

Any CHD

- CRT may be considered for patients with a severe subpulmonary RV dysfunction and dilatation despite interventions to decrease RV volume overload, NYHA function Class II—ambulatory IV and wide QRS complex ≥ 150 ms due to a complete RBBB
- NYHA function Class IV and severe ventricular dysfunction who would otherwise be candidates for heart transplantation or mechanical circulatory support

NOT Indicated for CRT in Adult Congenital Heart Disease

- Patients whose co-morbidities and/or frailty limit survival with good functional capacity to less than 1 year

INDICATIONS FOR CRT AS THE APPROPRIATE PACING MODALITY IN SPECIAL SITUATIONS WITH < 3 MONTHS OF GDMT^{5, 12, 13}

Criteria are met for a non-elective implantable cardioverter defibrillator (ICD) or pacemaker and based upon the low likelihood of improvement in symptoms and adequate recovery of LVEF, despite less than 3 months GDMT for heart failure or < 40 days post myocardial infarction or 3 months post revascularization, criteria for CRT are otherwise met. This avoids a second implantation procedure within less than 3 months.

BACKGROUND^{1, 3-5, 8}

CRT, which paces the left and right ventricle in rapid sequence, also known as biventricular pacing, improves coordination of ventricular contraction in the presence of a wide QRS complex in systolic heart failure.

CRT improves cardiac function and quality of life, and it decreases cardiac events and mortality among appropriately chosen patients. In the proper patient population, improved survival in patients with CRT can be greater than that provided by ICD insertion alone.

Guiding principles in the consideration of CRT:

- NYHA class is an important qualifying factor, with candidacy based on functional class, EF, and QRS duration.
- Bundle branch block or intraventricular conduction delay should be persistent, not rate related.⁵
- GDMT should have been in place continuously for at least 3 months^{3, 4, 8} and recovery of LVEF from myocardial infarction (40 days) if no intervening revascularization or > 3 months if revascularization was performed. Reversible causes (e.g., ischemia) should be excluded.
- The patient should have expected survival with reasonably good functional status for more than 1 year.^{3, 4, 10}

OVERVIEW

NYHA Class Definitions^{5, 14}

- Class I: No limitation of functional activity. Ordinary physical activity does not cause symptoms of HF
- Class II: Slight limitation of activity. Comfortable at rest but ordinary physical activity results in symptoms of HF
- Class III: Marked limitation of activity. Comfortable at rest but less than ordinary activity causes symptoms of HF
- Class IV: Unable to continue any physical activity without symptoms of HF, or symptoms of HF at rest

Heart Block Definitions³

- First Degree: All atrial beats are conducted to the ventricles, but with a delay of > 200 ms.
- Second Degree: Intermittent failure of conduction of single beats from atrium to ventricles.
 - Type I: Conducted beats have variable conduction times from atrium to ventricles.
 - Type II: Conducted beats have uniform conduction times from atrium to ventricles.
 - Advanced: Two or more consecutive non-conducted beats (premature atrial beats might not normally be conducted).
- Third Degree: No atrial beats are conducted from atrium to ventricle.

Guideline-Directed (or Optimal) Medical Therapy in Heart Failure⁸


- Angiotensin converting enzyme inhibitor (ACE-I), angiotensin receptor blocker (ARB), or combined angiotensin receptor inhibitor and neprilysin inhibitor (ARNI)
- Beta blocker

Other options/considerations for GDMT

- Addition of loop diuretic for all NYHA class II – IV patients
- Addition of hydralazine and nitrate for persistently symptomatic African Americans, NYHA class III-IV
- Addition of an aldosterone antagonist, provided eGFR is ≥ 30 ml/min/1.73m² and K⁺ < 5.0, NYHA class II-IV
- Not required for consideration of CRT: Ivabradine for NYHA class II – III, when a beta blocker has failed to reduce a sinus rate to < 70 bpm.

Algorithm for CRT Indications in Patients with Cardiomyopathy or HFrEF chart¹⁵

FIGURE 8 Algorithm for CRT Indications in Patients With Cardiomyopathy or HFrEF



Abbreviations

ACE-I	Angiotensin converting enzyme inhibitor
ARB	Angiotensin receptor blocker
ARNI	Combined angiotensin receptor inhibitor and neprilysin inhibitor
AV	Atrioventricular
CAD	Coronary artery disease, same as ischemic heart disease
CHD	Congenital heart disease
CHF	Congestive heart failure
CRT	Cardiac resynchronization therapy (also known as biventricular pacing)
CRT-D	Cardiac resynchronization therapy defibrillator
ECG	Electrocardiogram
EF	Ejection Fraction
eGFR	Estimated glomerular filtration rate
EPS	Electrophysiologic Study
GDMT	Guideline-Directed Medical Therapy
HCM	Hypertrophic Cardiomyopathy
HF	Heart failure
HFrEF	Heart failure with reduced ejection fraction
HV	His-ventricular
ICD	Implantable cardioverter-defibrillator
LBBB	Left bundle branch block
LV	Left ventricular/left ventricle
LVEF	Left ventricular ejection fraction
MI	Myocardial infarction
ms	Milliseconds
NYHA	New York Heart Association
RBBB	Right bundle branch block
RV	Right ventricle
SND	Sinus node dysfunction
SR	Sinus rhythm
STEMI	ST-Elevation Myocardial Infarction
VT	Ventricular tachycardia

REFERENCES

1. Brignole M, Auricchio A, Baron-Esquivias G, et al. 2013 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy: the Task Force on cardiac pacing and resynchronization therapy of the European Society of Cardiology (ESC). Developed in collaboration with the European Heart Rhythm Association (EHRA). *Eur Heart J*. Aug 2013;34(29):2281-329. doi:10.1093/eurheartj/eh150
2. Cleland JG, Daubert JC, Erdmann E, et al. The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med*. Apr 14 2005;352(15):1539-49. doi:10.1056/NEJMoa050496
3. Epstein AE, DiMarco JP, Ellenbogen KA, et al. 2012 ACCF/AHA/HRS focused update incorporated into the ACCF/AHA/HRS 2008 guidelines for device-based therapy of cardiac rhythm abnormalities: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *Circulation*. Jan 22 2013;127(3):e283-352. doi:10.1161/CIR.0b013e318276ce9b
4. Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. Jul 14 2016;37(27):2129-2200. doi:10.1093/eurheartj/ehw128
5. Russo AM, Stainback RF, Bailey SR, et al. ACCF/HRS/AHA/ASE/HFSA/SCAI/SCCT/SCMR 2013 appropriate use criteria for implantable cardioverter-defibrillators and cardiac resynchronization therapy: a report of the American College of Cardiology Foundation appropriate use criteria task force, Heart Rhythm Society, American Heart Association, American Society of Echocardiography, Heart Failure Society of America, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, and Society for Cardiovascular Magnetic Resonance. *J Am Coll Cardiol*. Mar 26 2013;61(12):1318-68. doi:10.1016/j.jacc.2012.12.017
6. Glikson M, Nielsen JC, Kronborg MB, et al. 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy: Developed by the Task Force on cardiac pacing and cardiac resynchronization therapy of the European Society of Cardiology (ESC) With the special contribution of the European Heart Rhythm Association (EHRA). *European Heart Journal*. 2021;42(35):3427-3520. doi:10.1093/eurheartj/ehab364
7. Baumgartner H, De Backer J, Babu-Narayan SV, et al. 2020 ESC Guidelines for the management of adult congenital heart disease. *Eur Heart J*. Feb 11 2021;42(6):563-645. doi:10.1093/eurheartj/ehaa554
8. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. May 3 2022;145(18):e876-e894. doi:10.1161/cir.0000000000001062
9. Hernández-Madrid A, Paul T, Abrams D, et al. Arrhythmias in congenital heart disease: a position paper of the European Heart Rhythm Association (EHRA), Association for European

Paediatric and Congenital Cardiology (AEPC), and the European Society of Cardiology (ESC) Working Group on Grown-up Congenital heart disease, endorsed by HRS, PACES, APHRS, and SOLAECE. *Europace*. Nov 1 2018;20(11):1719-1753. doi:10.1093/europace/eux380

10. Khairy P, Van Hare GF, Balaji S, et al. PACES/HRS Expert Consensus Statement on the Recognition and Management of Arrhythmias in Adult Congenital Heart Disease: developed in partnership between the Pediatric and Congenital Electrophysiology Society (PACES) and the Heart Rhythm Society (HRS). Endorsed by the governing bodies of PACES, HRS, the American College of Cardiology (ACC), the American Heart Association (AHA), the European Heart Rhythm Association (EHRA), the Canadian Heart Rhythm Society (CHRS), and the International Society for Adult Congenital Heart Disease (ISACHD). *Heart Rhythm*. Oct 2014;11(10):e102-65. doi:10.1016/j.hrthm.2014.05.009

11. Stout KK, Daniels CJ, Abouhosn JA, et al. 2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. Apr 2 2019;73(12):1494-1563. doi:10.1016/j.jacc.2018.08.1028

12. Kusumoto FM, Calkins H, Boehmer J, et al. HRS/ACC/AHA expert consensus statement on the use of implantable cardioverter-defibrillator therapy in patients who are not included or not well represented in clinical trials. *Heart Rhythm*. Jul 2014;11(7):1271-303. doi:10.1016/j.hrthm.2014.03.041

13. Marine JE, Russo AM. Primary prevention of sudden cardiac death in patients with cardiomyopathy and heart failure with reduced LVEF. Wolter Kluwer. Updated December 20, 2022. Accessed January 27, 2022. <https://www.uptodate.com/contents/primary-prevention-of-sudden-cardiac-death-in-patients-with-cardiomyopathy-and-heart-failure-with-reduced-lvef>

14. Goldman L, Hashimoto B, Cook EF, Loscalzo A. Comparative reproducibility and validity of systems for assessing cardiovascular functional class: advantages of a new specific activity scale. *Circulation*. Dec 1981;64(6):1227-34. doi:10.1161/01.cir.64.6.1227

15. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure. *Journal of the American College of Cardiology*. 2022;79(17):e263-e421. doi:doi:10.1016/j.jacc.2021.12.012

POLICY HISTORY

Date	Summary
April 2023	<ul style="list-style-type: none">• Added additional statement on atrial fibrillation• Added statement on ATTR• Added additional contraindication for patients with LVAD• Removed indication for Class I and CRT• Combined Class II- IV indications• Removed EF value for requirement for pacemaker• Added statement on clinical indications not addressed in this guideline
February 2022	<ul style="list-style-type: none">• Added blood stream infection and reversibility as contraindication• Reworded NYHA• Removed single ventricle and RV

Reviewed / Approved by Clinical Guideline Committee

Disclaimer: *Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as required by applicable provider agreements and laws or regulations. **Members should contact their Plan customer service representative for specific coverage information.***